




Alzheimer's Disease
Research Center



HEALTHY BRAIN AGING LECTURE SERIES: HEALTHY BRAIN AGING

Dr. DeCarli will discuss changes to the brain as we age and in the presence of common, treatable medical illnesses. By the end of the talk, you will be provided with actionable insights to optimize your brain health and enhance your quality of life.

September 5th, 2024, 6 p.m. - 7:30 p.m.

Education Building Room 1222

4610 X Street, Sacramento, CA 95817

Sponsored by:



**SUSAN & CHARLES
BERGHOFF foundation**

We envision a dementia-free world™



Charles DeCarli, M.D., FAAN, FAHA

Distinguished Professor of Neurology
Co-Director, Alzheimer's Disease Research Center
Department of Neurology and Center for Neuroscience

Thank you to Our Sponsor



SUSAN & CHARLES BERGHOFF *foundation*

We envision a dementia-free world™

Life's Essential 8 What You Can Do To Improve Your Brain Health

Charles DeCarli, MD
Victor and Genevieve
Orsi Chair in
Alzheimer's Research
Co-Director University
of California at Davis,
Alzheimer's Disease
Center



Dementia prevention, intervention, and care: 2020 report of the Lancet Commission



Gill Livingston, Jonathan Huntley, Andrew Sommerlad, David Ames, Clive Ballard, Sube Banerjee, Carol Brayne, Alistair Burns, Jiska Cohen-Mansfield, Claudia Cooper, Sergi G Costafreda, Amit Dias, Nick Fox, Laura N Gitlin, Robert Howard, Helen C Kales, Mika Kivimäki, Eric B Larson, Adesola Ogunniyi, Vasiliki Orgeta, Karen Ritchie, Kenneth Rockwood, Elizabeth L Sampson, Quincy Samus, Lon S Schneider, Geir Selbæk, Linda Teri, Naaheed Mukadam

Panel: Recommended strategies for dementia risk reduction


Risks are particularly high in more socially disadvantaged populations including in Black, Asian, and minority ethnic groups.

Population-wide

- Prioritise childhood education for all, worldwide
- Implement social public health policies that reduce hypertension risk in the entire population
- Develop policies that encourage social, cognitive, and physical activity across the life course for all (with no evidence for any specific activities being more protective)
- Scrutinise the risks for hearing loss throughout the life course, to reduce the risk of exposure to this risk factor
- Reduce the risk of serious brain trauma in relevant settings, including occupational and transport
- National and international policies to reduce population exposure to air pollution
- Continue to strengthen national and international efforts to reduce exposure to smoking, both for children and adults, and to reduce uptake and encourage cessation

Targeted on individuals

- Treat hypertension and aim for systolic blood pressure <130 mm Hg in midlife
- Use hearing aids for hearing loss; we need to help people wear hearing aids as many find them unacceptable, too difficult to use, or ineffective
- Avoid or discourage drinking 21 or more units of alcohol per week
- Prevent head trauma where an individual is at high risk
- Stopping smoking is beneficial regardless of age
- Reduce obesity and the linked condition of diabetes by healthy food availability and an environment to increase movement
- Sustain midlife, and possibly late-life physical activity



Risk score for the prediction of dementia risk in 20 years among middle aged people: a longitudinal, population-based study

Miia Kivipelto, Tiia Ngandu, Tiina Laatikainen, Bengt Winblad, Hilka Soininen, Jaakko Tuomilehto


Lancet Neurol 2006; 5: 735–41

Midlife risk score for the prediction of dementia four decades later


Lieza G. Exalto^{a,b}, Charles P. Quesenberry^a, Deborah Barnes^c, Miia Kivipelto^d,
Geert Jan Biessels^a, Rachel A. Whitmer^{b,*}


Alzheimer's & Dementia 10 (2014) 562–570

Step 1		Step 4		Step 2		Step 5		Step 3		Step 6		Predicted 40-year risk of dementia	
Age, y	Points	Cholesterol, mg/dL	Points	Education, y	Points	BMI, kg/m ²	Points	Sex	Points	Systolic blood Pressure, mm/Hg	Points	Total points	40-Year risk, %
40–46	0	<251	0	0–6	3	<30	0	Men	1	<140	0	0–1	10
47–53	3	≥251	2	7–9	2	≥30	2	Female	0	>140	2	2	11
54–55	4			>9	0							3	15
												4	17
												5	20
												6	21
												7	25
												8–14	29




Age-Specific Prevalence of Vascular Disease and Brain Volume






Future Risk of Stroke or Dementia at Age 65

Seshadri, S. et al.
Stroke 2006;37:345-350




Increasing odds of Dementia with Number of Vascular Risk Factors


Whitmer, et al, Neurology, 2005
~74% Caucasian




Luchsinger, et al, Neurology 2005
~23% Caucasian



Pathology of Dementia




Vascular Pathology of Dementia




Summary

- Vascular risk factors and consequential vascular disease is common to the aging process
- Vascular risk factors also increase risk for dementia
- Vascular pathology commonly accompanies Alzheimer's disease in dementia


IMPROVING COGNITIVE AGING





NHANES, National Health and Nutrition Examination Survey; NHES, National Health Examination Survey

Source: Burt VL, et al. Trends in the prevalence, awareness, treatment, and control of hypertension in the adult US population. Data from the health examination surveys, 1960 to 1991. Erratum in: Hypertension 1996;7(5):1192.




Reduction in SBP mmHg	% Reduction in Mortality		
	Stroke	CHD	Total
2	-6	-4	-3
3	-8	-5	-4
5	-14	-9	-7

BP, blood pressure; CHD, coronary heart disease; SBP, systolic blood pressure


Source: Whelton PK, et al. Primary prevention of hypertension: Clinical and public health advisory from The National High Blood Pressure Education Program. JAMA 2002;288:1882–8.

Trajectory of mortality rates from CHD and stroke, rate of uncontrolled high blood pressure, and prevalence of high blood cholesterol from 2004 to 2008.



Donald M. Lloyd-Jones et al. *Circulation*. 2010;121:586-613

Prevalence of Heart Disease and Stroke for those over 20 years of age




The projected effect of risk factor reduction on Alzheimer's disease prevalence



Deborah E Barnes, Kristine Yaffe

www.thelancet.com/neurology Vol 10 September 2011



Defining and Setting National Goals for Cardiovascular Health Promotion and Disease Reduction

The American Heart Association's Strategic Impact Goal Through 2020 and Beyond

Donald M. Lloyd-Jones, Yuling Hong, Darwin Labarthe, Dariush Mozaffarian, Lawrence J. Appel, Linda Van Horn, Kurt Greenlund, Stephen Daniels, Graham Nichol, Gordon F. Tomaselli, Donna K. Arnett, Gregg C. Fonarow, P. Michael Ho, Michael S. Lauer, Frederick A. Masoudi, Rose Marie Robertson, Véronique Roger, Lee H. Schwamm, Paul Sorlie, Clyde W. Yancy, Wayne D. Rosamond

and on behalf of the American Heart Association Strategic Planning Task Force and Statistics Committee


Circulation. 2010;121:586-613

Component	Ideal (2 Points)	Intermediate (1 Point)	Poor (0 Points)
Smoking	Never or former >1 year	Former \leq 1 year	Current
Healthy diet score*	4 to 5 points	2 to 3 points	0 to 1 points
Physical activity [†]	\geq 4 bouts per week of intense physical activity sufficient to work up a sweat	1 to 3 bouts per week of intense physical activity sufficient to work up a sweat	No intense physical activity sufficient to work up a sweat
Body mass index	$<$ 25 kg/m ²	25 to 29.9 kg/m ²	\geq 30 kg/m ²
Blood pressure	$<$ 120/ $<$ 80 mm Hg untreated	SBP 120 to 139 or DBP 80 to 89 mm Hg or treated to ideal level	SBP \geq 140 or DBP \geq 90 mm Hg
Total cholesterol	$<$ 5.18 mmol/L ($<$ 200 mg/dL) untreated	5.18 to 6.19 mmol/L (200 to 239 mg/dL) or treated to ideal level	\geq 6.22 mmol/L (\geq 240 mg/dL)
Fasting glucose	$<$ 5.55 mmol/L ($<$ 100 mg/dL) untreated	5.55 to 6.94 mmol/L (100 to 125 mg/dL) or treated to ideal level	\geq 6.99 mmol/L (\geq 126 mg/dL)

Life's Essential 8: Updating and Enhancing the American Heart Association's Construct of Cardiovascular Health: A Presidential Advisory From the American Heart Association

Donald M. Lloyd-Jones, MD, ScM, FAHA, Chair; Norrina B. Allen, PhD, MPH, FAHA; Cheryl A.M. Anderson, PhD, MPH, MS, FAHA; Terrie Black, DNP, MBA, CRRN, FAHA; LaPrincess C. Brewer, MD, MPH; Randi E. Foraker, PhD, MA, FAHA; Michael A. Grandner, PhD, MTR, FAHA; Helen Lavretsky, MD, MS; Amanda Marma Perak, MD, MS, FAHA; Garima Sharma, MD; Wayne Rosamond, PhD, MS, FAHA; on behalf of the American Heart Association





A Few Words About Diet



Dietary habits show strongest evidence for causal effects on cardiovascular events, diabetes, and/or obesity



Recommend dietary pattern based on foods rather than nutrients



Inclusion of as few as possible elements with minimal overlap with each other while at the same time having some overlap with other relevant dietary guidelines

Dietary Approaches to Stop Hypertension (DASH)



Fruits and vegetables:
 ≥ 4.5 cups per day



Fish: \geq two 3.5-oz servings per week (preferably oily fish)



Fiber-rich whole grains: \geq three 1-oz-equivalent servings per day



Sodium: < 1500 mg per day



Sugar-sweetened beverages: ≤ 450 kcal (36 oz) per week

Secondary Dietary Metrics


Nuts, legumes, and seeds:
 ≥ 4 servings per week

Processed meats: none or
 ≤ 2 servings per week

Saturated fat: $< 7\%$ of total
energy intake



Other Diets



THE MIND DIET

- Green, leafy vegetables – daily 
- Other vegetables – daily 
- Nuts – at least once a day 
- Berries – at least twice a week 
- Beans or legumes – at least every other day 
- Whole grains – at least 3 servings a day 
- Fish – at least once a week 
- Poultry – at least twice a week 

A Few Words About Sleep

It is not all about the length!

- Too short is bad
- Too long is bad


Its about the quality!

- Restful, deep sleep is best
- Awaking spontaneously is best

Its about the oxygen!

- Sleep apnea reduces oxygen to the brain
 - Repeated awakening
 - Day time sleepiness

- Loud snoring
- Episodes in which you stop breathing during sleep — which would be reported by another person
- Gasping for air during sleep
- Awakening with a dry mouth
- Morning headache
- Difficulty staying asleep (insomnia)
- Excessive daytime sleepiness (hypersomnia)
- Difficulty paying attention while awake
- Irritability



Risk Factors for Sleep Apnea

- **Excess weight:** Fat deposits around your upper airway can obstruct your breathing
- **Neck circumference:** People with thicker necks might have narrower airways
- **A narrowed airway:** Tonsils or adenoids also can enlarge and block the airway, particularly in children.
- **Being male**
- **Being older**
- **Positive Family history**

Risk Factors for Sleep Apnea (cont'd)



USE OF ALCOHOL, SEDATIVES OR TRANQUILIZERS.

THESE SUBSTANCES RELAX THE MUSCLES IN YOUR THROAT, WHICH CAN WORSEN OBSTRUCTIVE SLEEP APNEA



SMOKING



NASAL CONGESTION



MEDICAL CONDITIONS:
CONGESTIVE HEART FAILURE, HIGH BLOOD PRESSURE, TYPE 2 DIABETES AND LUNG DISEASES

Monitoring your Cardiovascular Health



American Heart Association.
My Life Check®


**GOOD HABITS BUILD
BETTER HEALTH**

We've helped millions of people make healthier choices.


The AHA is the nation's oldest and largest voluntary organization dedicated to fighting heart disease and stroke. For nearly 100 years, we've been helping people like you live longer, healthier lives.

<https://mlc.heart.org/>


Overall Cardiovascular Health (CVH)



Health Behaviors




Health Factors



The American Heart Association Life's Simple 7 and Incident Cognitive Impairment: The REasons for Geographic And Racial Differences in Stroke (REGARDS) Study


Evan L. Thacker, PhD; Sarah R. Gillett, PhD; Virginia G. Wadley, PhD; Frederick W. Unverzagt, PhD; Suzanne E. Judd, PhD; Leslie A. McClure, PhD; Virginia J. Howard, PhD; Mary Cushman, MD, MSc

J Am Heart Assoc. 2014



Study Design

- 17,761 Individuals > 45 years of age
 - Free of Stroke and Dementia
- Study Duration: 2003-2012
- Biennial Assessment of Cognition
 - Word list immediate and delayed recall
 - Animal fluency



Life's Simple 7 Component	Adjusted Odds Ratio* (95% CI)
Smoking	
Poor	1.00 (reference)
Intermediate or ideal [†]	0.77 (0.60, 0.98)
Healthy diet score	
Poor	1.00 (reference)
Intermediate or ideal [†]	0.90 (0.68, 1.18)
Physical activity	
Poor	1.00 (reference)
Intermediate	0.88 (0.72, 1.08)
Ideal	0.94 (0.76, 1.17)
Body mass index	
Poor	1.00 (reference)
Intermediate	0.77 (0.63, 0.95)
Ideal	0.71 (0.56, 0.91)
Blood pressure	
Poor	1.00 (reference)
Intermediate	0.79 (0.64, 0.97)
Ideal	0.86 (0.65, 1.14)
Total cholesterol	
Poor	1.00 (reference)
Intermediate	1.08 (0.80, 1.45)
Ideal	1.04 (0.77, 1.40)
Fasting glucose	
Poor	1.00 (reference)
Intermediate	0.79 (0.59, 1.07)
Ideal	0.72 (0.53, 0.97)

Impact of Individual Components

Association of Ideal Cardiovascular Health With Vascular Brain Injury and Incident Dementia


Matthew P. Pase, PhD; Alexa Beiser, PhD; Danielle Enserro, MA;
 Vanessa Xanthakis, PhD; Hugo Aparicio, MD; Claudia L. Satizabal, PhD;
 Jayandra J. Himali, PhD; Carlos S. Kase, MD; Ramachandran S. Vasan, MD;
 Charles DeCarli, MD; Sudha Seshadri, MD

Event	Remote Ideal CVH		Ideal CVH frequencies, n (%)	
	HR (95% CI)	P Value		
Stroke	0.79 (0.66–0.94)	0.01	0	22 (0.8)
			1	245 (9.3)
			2	638 (24.3)
			3	734 (27.9)
			4	569 (21.6)
			5	275 (10.5)
			6	124 (4.7)
			7	24 (0.9)
All-cause dementia	0.80 (0.67–0.97)	0.02		
Alzheimer disease	0.79 (0.64–0.98)	0.006		
Vascular dementia	0.61 (0.39–0.95)	0.03		

Measures	Recent Ideal CVH		Remote Ideal CVH	
	$\beta \pm SE$	P Value	$\beta \pm SE$	P Value
Cognitive decline				
Global decline	0.003±0.002	0.07	0.006±0.002	0.002
Visual reproductions delayed	0.02±0.01	0.01	0.02±0.01	0.01
Similarities	0.02±0.01	0.04	0.04±0.01	<0.001
Trail A	0.001±0.002	0.46	-0.002±0.002	0.43
Trail B	-0.01±0.004	0.13	-0.01±0.004	0.08
Logical Memory Delayed	-0.01±0.01	0.51	0.01±0.01	0.44
Brain atrophy and white-matter injury				
Total brain volume	0.09±0.08	0.26	0.19±0.08	0.02
Frontal brain volume	0.31±0.10	0.003	0.15±0.11	0.16
Lateral ventricular volume	0.02±0.01	0.10	0.002±0.02	0.88
WMHV	-0.0002±0.01	0.98	0.0003±0.01	0.97

RESEARCH ARTICLE

Cardiovascular health and cognitive outcomes: Findings from a biracial population-based study in the United States

Anisa Dhana^{1,2}  | Charles S. DeCarli³ | Klodian Dhana^{1,2} | Pankaja Desai^{1,2} |
Thomas M. Holland^{1,2} | Denis A. Evans^{1,2} | Kumar B. Rajan^{1,2,3}

Journal of Alzheimer's Disease 53 (2016) 955–965
DOI 10.3233/JAD-151125
IOS Press


955

Life's Simple 7's Cardiovascular Health Metrics are Associated with Hispanic/Latino Neurocognitive Function: HCHS/SOL Results

Hector M. González^{a,*}, Wassim Tarrat^b, Natalia Gouskova^c, Carlos J. Rodríguez^d, Tatjana Rundek^e, Ellen Grober^f, Amber Pirzada^g, Patricia González^h, Pamela L. Lutseyⁱ, Alvaro Camacho^j, Martha L. Daviglus^g, Clinton Wright^e and Thomas H. Mosley^k

Physical Activity, Brain Volume, and Dementia Risk: The Framingham Study

Zaldy S. Tan,^{1,2,*} Nicole L. Spartano,^{2,3,*} Alexa S. Beiser,^{2,4,5} Charles DeCarli,⁶ Sanford H. Auerbach,^{2,4} Ramachandran S. Vasan,^{2,3} and Sudha Seshadri²




	$\beta \pm SE$	
	Age and Sex Adjusted	<i>p</i> Value
TCBV		
Per <i>SD</i>	0.24 ± 0.06	<.001
Q1	1.00 (Referent)	
Q2	0.02 ± 0.20	.914
Q3	0.26 ± 0.20	.191
Q4	0.56 ± 0.20	.005
Q5	0.54 ± 0.20	.007
HPV		
Per <i>SD</i>	0.004 ± 0.001	.003
Q1	1.00 (Referent)	
Q2	0.001 ± 0.004	.713
Q3	0.001 ± 0.004	.746
Q4	0.006 ± 0.004	.107
Q5	0.010 ± 0.004	.007

AHA/ASA Presidential Advisory

Defining Optimal Brain Health in Adults A Presidential Advisory From the American Heart Association/ American Stroke Association

Philip B. Gorelick, MD, MPH, FAHA, Chair*; Karen L. Furie, MD, MPH, FAHA, Co-Chair†; Costantino Iadecola, MD, FAHA, Co-Chair†; Eric E. Smith, MD, MPH, FAHA‡; Salina P. Waddy, MD§; Donald M. Lloyd-Jones, MD, ScM, FAHA||; Hee-Joon Bae, MD, PhD, FAHA; Mary Ann Bauman, MD; Martin Dichgans, MD; Pamela W. Duncan, PhD, PT, FAHA; Meighan Girgus; Virginia J. Howard, PhD, FAHA; Ronald M. Lazar, PhD, FAHA; Sudha Seshadri, MD, FAHA; Fernando D. Testai, MD, PhD, MS, FAHA; Stephen van Gaal, MD; Kristine Yaffe, MD, FAHA; Hank Wasiak, MBA; Charlotte Zerna, MD, MSc; on behalf of the American Heart Association/
American Stroke Association


Stroke. 2017;48:e284-e303



Optimal Brain Health




Cognitive Dysfunction




Individual ¹⁶⁸
Check health status with AHA's Life's Simple 7 (http://www.heart.org)
Remain physically active
Eat a healthy diet; evidence suggests that a Mediterranean-style diet preserves cognitive function better than a low-fat diet
Address vascular risk factors, if present, with a primary care practitioner
Pursue cognitively stimulating and rewarding activities
Address mental health concerns with a primary care practitioner or specialist as needed
Healthcare practitioners
Apply primordial and primary preventive care for cardiovascular disease and stroke according to AHA/ASA guidelines ^{9,124,142,163,164}
Diagnose and treat symptomatic stroke according to AHA/ASA guidelines ¹⁶⁵⁻¹⁶⁷
Administer brief screens to monitor cognitive status

Recommendations for Promotion and Maintenance of Optimal Brain Health

Health systems
Support patients by providing access to preventive care and lifestyle modification
Support good-quality care for stroke ¹⁶⁹ and for primary prevention of cardiovascular disease ¹⁷⁰
Public health, health policy, private sector ^{9,168}
Disseminate knowledge of potentially modifiable risk factors for cognitive decline and dementia
Provide tools and resources to maintain healthy lifestyles such as the AHA Healthy for Good program ¹⁷¹
Provide opportunities for stimulating cognitive, physical, and social activities
Maintain a healthy environment, including neighborhoods that promote cognitive and physical activity
Fund research on risk factors for cognitive decline and dementia and how to intervene to reduce risk



IMPROVING COGNITIVE AGING



Conclusions



Treatment of vascular risk factors and promotion of “vascular health” is likely to have a strong public health benefit to reduce late-life dementia



Current efforts to improve vascular health with “Life’s Essential 8” provide a public health opportunity to assess the efficacy of this approach

Life's Essential 8



DON'T SMOKE

Life's Essential 8



DON'T SMOKE



WATCH YOUR
WEIGHT

Life's Essential 8



DON'T SMOKE



WATCH YOUR
WEIGHT



EAT A HEALTHY
DIET

Life's Essential 8



DON'T SMOKE



WATCH YOUR
WEIGHT



EAT A HEALTHY
DIET



EXERCISE

Life's Essential 8



DON'T SMOKE



WATCH YOUR
WEIGHT



EAT A HEALTHY
DIET



EXERCISE



CONTROL
YOUR BLOOD
PRESSURE

Life's Essential 8



DON'T SMOKE



WATCH YOUR
WEIGHT




EAT A HEALTHY
DIET



EXERCISE



CONTROL
YOUR BLOOD
PRESSURE



CONTROL
YOUR BLOOD
SUGAR

Life's Essential 8



DON'T SMOKE



WATCH YOUR
WEIGHT




EAT A HEALTHY
DIET



EXERCISE



CONTROL
YOUR BLOOD
PRESSURE



CONTROL
YOUR BLOOD
SUGAR



CONTROL
YOUR BLOOD
CHOLESTEROL

Life's Essential 8



DON'T SMOKE



WATCH YOUR
WEIGHT




EAT A HEALTHY
DIET



EXERCISE



CONTROL
YOUR BLOOD
PRESSURE



CONTROL
YOUR BLOOD
SUGAR



CONTROL
YOUR BLOOD
CHOLESTEROL



SLEEP WELL

Life's Essential 8

- Don't Smoke
- Watch Your Weight
- Eat A Healthy Diet
- Exercise
- Control Your Blood Pressure
- Control Your Blood Sugar
- Control Your Blood Cholesterol
- Sleep Well



Ways to Get Involved

- Join the **Research Registry**
- **Make a donation** to fund research and community education programs
- **Attend our other workshops**, such as the Caregiver Bootcamp
- Join us at the next **Healthy Brain Aging Lectures**
 - Thursday, October 3 – **US Pointer Study** with Dr. Rachel Whitmer
 - Thursday, November 7 – **Tools and Strategies to Support Memory Loss** with Dr. Alyssa Weakley



Alzheimer's Disease
Research Center

Thank you for attending the Healthy Brain Aging Lecture!



Charles DeCarli, M.D., FAAN, FAHA

Distinguished Professor of Neurology
Co-Director, Alzheimer's Disease Research Center
Department of Neurology and Center for Neuroscience

Please complete the
evaluation by scanning the
QR Code or visit
<https://tinyurl.com/yhyn8p7z>

