

# Fluoroscopic Swallow Study Predictors of Tracheoesophageal Puncture Voice Quality

Kristiana M. Jordan, BA<sup>1</sup>; Nina W. Zhao, MD, MAEd<sup>2</sup>; Lisa M. Evangelista, CScD, CCC-SLP, BCS-S<sup>2</sup>; Tess T. Andrews, MS, CCC-SLP<sup>2</sup>; Leia K. Chapman, MS, CCC-SLP<sup>2</sup>; Angela A. Colback, BA<sup>1</sup>; Peter C. Belafsky, MD, PhD<sup>4</sup>; Maggie Kuhn, MD, MAS<sup>4</sup>  
<sup>1</sup>University of California, Davis School of Medicine; <sup>2</sup>Department of Otolaryngology – Head and Neck Surgery, University of California, Davis

## OBJECTIVE

Tracheoesophageal puncture (TEP) is the gold standard for voice rehabilitation after total laryngectomy (TL). TEP voicing can be challenging & unpredictable. The dynamic swallow study is a fluoroscopic swallow study part of standard clinical care prior to TEP placement.





Figure 1: Pre-laryngectomy fluoroscopic study demonstrating mild dysphagia consistent with late radiation effects characterized by mild residue in the vallecula and pyriform sinus. Biomechanical impairments includes narrowing of the PES segment at the level of C5-C6.

Our study aimed to determine which **objective dynamic swallow study (DSS) parameters predict improved voice outcomes** in patients who have TEP placement.

## METHODS



## RESULTS

N=14 patients had full data set for pre- and post-laryngectomy DSS parameters and SLP-voice ratings:

Patient Characteristic	Global Voice Rating		
	Below Average	Average	Above Average
<b>N=14 (total)</b>			
Age			
43-70 years (N=7)	43%	43%	43%
70+ years (N=7)	43%	14%	14%
Dysphagia			
Yes (N=13)	46%	31%	23%
No (N=1)	0%	100%	0%
History of acid reflux (Y/N)			
Yes (N=2)	0%	100%	0%
No (N=10)	40%	30%	30%
N/A (N=2)	100%	0%	0%
Complications since TEP placement			
Yes (N=5)	60%	20%	20%
No (N=9)	33%	44%	22%

Table 1: Breakdown of demographic information and clinical parameters collected. Age distribution is split evenly in each age group, with most patients experiencing dysphagia. Below average = global TEP score of 0; average = global TEP score of 1; above average = global TEP score of 2.

N=14 (total)	Global Voice Rating		
	Below Average	Average	Above Average
<b>Comorbidities</b>			
Aspiration Pneumonia (N=2)	7%	7%	0%
GERD (N=1)	0%	7%	0%
Heart Disease (N=7)	21%	21%	7%
Hypertension (N=7)	21%	21%	7%
Hypothyroidism (N=4)	14%	14%	0%
Hyperlipidemia (N=2)	7%	7%	0%
Lung Disease (N=7)	29%	7%	14%
None (N=1)	0%	0%	7%

Table 2: Highlights the comorbidities found in the patient population examined. Most common comorbidities include hypertension and heart disease, primarily in patients with below average to average global TEP voice scores.




Figure 3: Multivariate linear regression demonstrated pre-laryngectomy fluoroscopy measures (A) pharyngoesophageal segment size (AP) ( $\beta = -1.6356$ ; [95% CI, 0.10-1.0.35];  $p < 0.05$ ), (B) pharyngeal constriction ratio ( $\beta = -1.6591$ ; [95% CI, 0.05-0.71];  $p < 0.05$ ), and (C) total pharyngeal transit time ( $\beta = -0.3957$ ; [95% CI, 0.51-0.89];  $p < 0.05$ ) to be independent predictors of global TEP voice ratings.  $F(6,5) = 19.68$   $p < 0.0025$   $R^2 = 0.9106$ .

## CONCLUSIONS

- We found the following pre-laryngectomy fluoroscopy parameters to be independent predictors of global TEP voicing:

### 1. Pharyngoesophageal segment size (AP)

- As PES size ↓, Global TEP Voice Rating ↓

### 2. Pharyngeal constriction ratio (PCR)

- As PCR ↓, Global TEP Voice Rating ↓

### 3. Total pharyngeal transit time

- As total pharyngeal transit time ↓, Global TEP Voice Rating ↓

- No independent predictors found in post-laryngectomy dynamic swallow study measurements.
- Future studies should explore prospective voice rating modalities for a more accurate voice rating score. Additionally, a larger sample size is necessary to further investigate the findings of this study.

## REFERENCES

- Zhang, Teng et al. "The relationship between biomechanics of pharyngoesophageal segment and tracheoesophageal phonation." *Scientific reports* vol. 9,1 9722. 5 Jul. 2019, doi:10.1038/s41598-019-46223-7
- Evangelista, Lisa et al. "Association of Functional Outcomes in Tracheoesophageal Voicing With Intratracheal Pressures and Esophagram Findings." *JAMA otolaryngology-- head & neck surgery* vol. 147,12 (2021): 1065-1070. doi:10.1001/jamaoto.2021.2409
- Takeshita-Monaretti, Telma Kioko et al. "Correlation of maximum phonation time and vocal intensity with intraluminal esophageal and pharyngoesophageal pressure in total laryngectomees." *The Annals of otology, rhinology, and laryngology* vol. 123,11 (2014): 811-6. doi:10.1177/0003489414538766