

Echocardiography, Targeted Neonatal Echocardiography (TNE) and Cardiac POCUS use in the NICU


Joint guideline developed in collaboration with Pediatric Cardiology, Neonatology, Neonatal Hemodynamics and POCUS Teams

A) Defining terms Echocardiography, TNE and Cardiac POCUS

1. **Echocardiography** refers to formal structural or functional echocardiography performed or read by the pediatric cardiologist.
2. **Targeted Neonatal Echocardiography** (TNE or TnECHO or functional echocardiography) refers to a limited echocardiography performed by a neonatologist formally trained in functional echocardiography, such as a comprehensive assessment or hemodynamic evaluation of patent ductus arteriosus (PDA), pulmonary hypertension (PH or PPHN), Hypoxic Ischemic Encephalopathy (HIE), or shock. This entails a structured assessment and evaluation using a standardized scanning protocol.
 - **First echocardiography, performed or read by a pediatric cardiologist**, should have a comprehensive structural assessment to rule out congenital heart defects (CHDs) and ideally it should be performed before TNE.
 - If the first echocardiography is performed by a TNE-trained person, a formal echocardiography should be obtained by cardiologists within 24 hours.
 - All TNE echo images should be stored and available for review. The findings and interpretation report should be documented in the EPIC.
 - All TNE evaluations should be performed by person trained in TNE who should make clinical decision in the clinical context and hemodynamic evaluation. When TnECHO performed by TNE trainee, then it should be for educational purpose only and should be reviewed by TNE expert to make a clinical decision.
3. **Cardiac POCUS** refers to a targeted, focused cardiac assessment to answer any clinical-specific question, such as qualitative assessment of cardiac function, detection of pericardial effusion or tamponade, detection of line position, and other urgent evaluations.

- **Cardiac POCUS** has a limited scope, with indications that are usually used in emergency situations or when a specific clinical question needs to be answered.
- Cardiac POCUS should not be used for screening of CHDs or comprehensive hemodynamic evaluation.
- If abnormality is detected on cardiac POCUS or the patient needs intervention, it should be promptly discussed with pediatric cardiology.

Echo, TNE & Cardiac POCUS - Difference?



- Different levels of training, indications and limitations And
- **Cardiac POCUS ≠ TNE (NPE) ≠ full echo**

Indications and Scope of Practice:

A). Formal echocardiography in the NICU: The following are the common indications of requesting a formal echocardiography in the NICU. In addition to these common indications, a formal echocardiography should be discussed with a pediatric cardiologist if there are any cardiac concerns by the attending neonatologist.

- Suspicion of CHDs including in infants of diabetic mothers, suspected trisomy 21 or syndromic association.
- Antenatal diagnosis or suspicion of CHDs.
- Diagnosis of PH and subsequent serial echocardiography as per PH or pediatric cardiology team.
- Subsequent echocardiography to guide management in infants with confirmed with CHDs.
- All cases with moderate to severe HIE needing therapeutic hypothermia with hemodynamic instability should have an echocardiography performed within 12 hours after birth, preferably soon after starting cooling.
- All cases with moderate to severe PH should have an echo performed as soon as possible after clinical diagnosis, especially if they need iNO or inotropic support.

- All cases with congenital diaphragmatic hernia (CDH) should have a formal echocardiography performed as soon as possible after birth.
- All extremely preterm infants <29 weeks of gestation should have a formal echocardiography done between 48-96 hours after birth.

B). TNE or TnECHO indications: Preferably, the 1st echo should be performed by pediatric cardiology before limited TNE assessments. When TNE is performed before a formal echocardiography, it should be obtained within 24 hours after performing TNE. The common indications of performing TNE include the following:

- **Diagnosis and hemodynamic evaluation of PDA** – all cases should have a formal echocardiography performed before medical treatment or intervention. Subsequent formal echocardiography should be obtained as requested by pediatric cardiology, before transcatheter device closure, surgical ligation, or clinical concerns from the neonatologist or TNE expert.
- **Diagnosis and hemodynamic evaluation of PPHN or acute pulmonary hypertension** - all cases should have a formal echocardiography as soon as possible after diagnosis or within 24 hours when 1st assessment is done on TNE. Subsequent formal echocardiography should be obtained as requested by pediatric cardiology, pulmonary hypertension team, or clinical concerns from the neonatologist or TNE expert.
- Diagnosis and hemodynamic evaluation of patients with shock or hypotension
- Diagnosis and hemodynamic evaluation of patients with HIE
- Qualitative or quantitative assessment of cardiac function
- Detection of pericardial effusion or cardiac tamponade
- Guide pericardiocentesis in an emergency

C). Cardiac POCUS indications: Targeted limited non-protocolized cardiac assessment for specific indication:

- Qualitative assessment of cardiac function / cardiac contractility
- Qualitative assessment of cardiac filling in an emergency
- Detection of cardiac tamponade or pericardial effusion
- Guide pericardiocentesis in an emergency
- Central line (PICC or UVC) tip position
- Limited assessment of pulmonary hypertension or ductal patency in emergency

Scope of TNE or cardiac POCUS assessment in children with known or suspected CHDs: TNE and cardiac POCUS may be utilized by trained neonatologists for rapid hemodynamic assessment in critically ill infants. However, for all patients with:

- Known or suspected congenital heart disease,
- Moderate or greater ventricular dysfunction, or

- Moderate or greater atrioventricular valve regurgitation,
- Moderate or greater ventricular hypertrophy
- Moderate or greater pericardial effusion

the first-line cardiac imaging must be a comprehensive formal echocardiogram performed and interpreted by pediatric cardiology.

If urgent imaging is needed for hemodynamic instability or uncertainty, pediatric cardiology should be contacted immediately to determine the most appropriate and time-sensitive diagnostic pathway and be actively involved in the management of the patient and subsequent imaging.

Escalation of Findings on TNE or cardiac POCUS assessment

Neonatologists should promptly escalate cardiac POCUS or TNE findings to the on-call pediatric cardiology attending if any of the following are identified:

- Unclear or concerning ductal or atrial level shunting
- Unexplained ventricular hypertrophy or decreased function
- Evidence of pulmonary hypertension unresponsive to initial therapy
- Structural abnormalities not previously documented

Continued Training, Quality Assurance, and Governance

- Joint quality review with pediatric cardiology - quarterly meetings to review images, discuss diagnostic variability, and ensure best practices as a QI measure.

References:

1. Singh Y, Tissot C, Fraga MV, Yousef N, Cortes RG, Lopez J, et al. International evidence-based guidelines on Point of Care Ultrasound (POCUS) for critically ill neonates and children issued by the POCUS Working Group of the European Society of Paediatric and Neonatal Intensive Care (ESPNIC). *Crit Care*. 2020 Feb 24;24(1):65. doi: 10.1186/s13054-020-2787-9. PMID: 32093763.
2. McNamara PJ, Jain A, El-Khuffash A, Giesinger R, Weisz D, Freud L, Levy PT, Bhombal S, de Boode W, Leone T, Richards B, Singh Y, Acevedo JM, Simpson J, Noori S, Lai WW. Guidelines and Recommendations for Targeted Neonatal Echocardiography and Cardiac Point-of-Care Ultrasound in the Neonatal Intensive Care Unit: An Update from the American Society of Echocardiography. *J Am Soc Echocardiogr*. 2024 Feb;37(2):171-215. doi:

- 10.1016/j.echo.2023.11.016. Erratum in: J Am Soc Echocardiogr. 2024 Jul;37(7):722. doi: 10.1016/j.echo.2024.04.006. PMID: 38309835.
3. Singh Y, Gupta S, Groves AM, Gandhi A, Thomson J, Qureshi S, Simpson JM. Expert consensus statement 'Neonatologist-performed Echocardiography (NoPE)-training and accreditation in UK. Eur J Pediatr. 2016 Feb;175(2):281-7. doi: 10.1007/s00431-015-2633-2. Epub 2015 Sep 12. PMID: 26362538.
 4. de Boode WP, Singh Y, Gupta S, Austin T, Bohlin K, Dempsey E, Groves A, Eriksen BH, van Laere D, Molnar Z, Nestaas E, Rogerson S, Schubert U, Tissot C, van der Lee R, van Overmeire B, El-Khuffash A. Recommendations for neonatologist performed echocardiography in Europe: Consensus Statement endorsed by European Society for Paediatric Research (ESPR) and European Society for Neonatology (ESN). Pediatr Res. 2016 Oct;80(4):465-71. doi: 10.1038/pr.2016.126. Epub 2016 Jun 8. PMID: 27384404.
 5. Fraga MV, Bhombal S, Juliano C, Kim M, Groves AM, Chan B, Singh Y; National Neonatal POCUS Collaborative. Neonatal point-of-care ultrasound-guidelines for training, credentialing and quality assurance. J Perinatol. 2025 Aug 1. doi: 10.1038/s41372-025-02367-1. Epub ahead of print. PMID: 40751094.
 6. Singh Y, Roehr CC, Tissot C, et al. European Special Interest Group 'Neonatologist Performed Echocardiography' (NPE). Education, training, and accreditation of Neonatologist Performed Echocardiography in Europe-framework for practice. Pediatr Res. 2018 Jul;84(Suppl 1):13-17. doi: 10.1038/s41390-018-0078-9. PMID: 30072809.
 7. Elsayed Y, Wahab MGA, Mohamed A, Fadel NB, Bhombal S, Yousef N, Fraga MV, Afifi J, Suryawanshi P, Hyderi A, Katheria A, Kluckow M, De Luca D, Singh Y. Point-of-care ultrasound (POCUS) protocol for systematic assessment of the crashing neonate-expert consensus statement of the international crashing neonate working group. Eur J Pediatr. 2023 Jan;182(1):53-66. doi: 10.1007/s00431-022-04636-z. Epub 2022 Oct 14. PMID: 36239816.
 8. Singh Y: Echocardiography in the neonatal unit: current status and future prospects. Expert Rev Med Devices. 2024 Apr;21(4):307-316. doi: 10.1080/17434440.2024.2334449. Epub 2024 Mar 25. PMID: 38526192.
 9. Chan B, Mnyavanu N, Bhombal S, Fraga MV, Groves AM, Marshall S, Mukthapuram S, Singh Y; members of National Neonatal POCUS Collaborative. Essentials of Point-of-Care Ultrasound Coding and Billing at the Neonatal Intensive Care Unit Setting in the United States. Am J Perinatol. 2024 May 2. doi: 10.1055/s-0044-1786721. Epub ahead of print. PMID: 38698594.
 10. Singh Y, Lakshminrusimha S. Perinatal cardiovascular physiology and recognition of critical congenital heart defects. Clinics in Perinatology 2021; 48: 573-594.

Reviewed by: Yogen Singh, MD; Sherzana Sunderji, MD; Frank Ing, MD; Jay Yeh, MD; Steve McElroy, MD; Belinda Chan, MD

Medical Legal Disclaimer:

Welcome to the UC Davis Health, Department of Pediatrics, Clinical Practice Guidelines Website. All health and health-related information contained within the Site is intended chiefly for use as a resource by the Department's clinical staff and trainees in the course and scope of their approved functions/activities (although it may be accessible by others via the internet). This Site is not intended to be used as a substitute for the exercise of independent professional judgment. These clinical pathways are intended to be a guide for practitioners and may need to be adapted for each specific patient based on the practitioner's professional judgment, consideration of any unique circumstances, the needs of each patient and their family, and/or the availability of various resources at the health care institution where the patient is located. Efforts are made to ensure that the material within this Site is accurate and timely but is provided without warranty for quality or accuracy. The Regents of the University of California; University of California, Davis; University of California, Davis, Health nor any other contributing author is responsible for any errors or omissions in any information provided or the results obtained from the use of such information. Some pages within this Site, for the convenience of users, are linked to or may refer to websites not managed by UC Davis Health. UC Davis Health does not control or take responsibility for the content of these websites, and the views and opinions of the documents in this Site do not imply endorsement or credibility of the service, information or product offered through the linked sites by UC Davis Health. UC Davis Health provides limited personal permission to use the Site. This Site is limited in that you may not:

- Use, download or print material from this site for commercial use such as selling, creating course packets, or posting information on another website.
- Change or delete propriety notices from material downloaded or printed from it. · Post or transmit any unlawful, threatening, libelous, defamatory, obscene, scandalous, inflammatory, pornographic, or profane material, any propriety information belonging to others or any material that could be deemed as or encourage criminal activity, give rise to civil liability, or otherwise violate the law.
- Use the Site in a manner contrary to any applicable law.

You should assume that everything you see or read on this Site is copyrighted by University of California or others unless otherwise noted. You may download information from this Site as long as it is not used for commercial purposes, and you retain the proprietary notices. You may not use, modify, make multiple copies, or distribute or transmit the contents of this Site for public or commercial purposes without the express consent of UC Davis Health.