

Adult Post-Operative Pain Management

Pain medication is given immediately following your surgery to help with post-operative pain. Upon your discharge home, we suggest scheduled doses of Acetaminophen (tylenol) every 6 hours and Ibuprofen (Motrin) every 6 hours, alternating between medications every 3 hours (i.e. Take tylenol and wait 3 hours, then take Motrin and wait 3 hours, repeat) for the first 3-4 days after surgery. If you are without significant pain, medications can be taken more infrequently. It is important to follow dosing instructions on the medication bottle or prescription.

If you need to call after clinic hours for a concern, call the UC Davis Medical Center hospital operator at 916-734-2011 and ask for the "physician on call for ENT."

UC Davis Otolaryngology
Facial Plastic and Reconstructive Surgery
2521 Stockton Blvd
Sacramento, CA 95817
916-734-5400

Sample of medication dosing schedule		
Give dose of:	Time:	Given:
Acetaminophen	12 a.m.	
Ibuprofen	3 a.m.	
Acetaminophen	6 a.m.	
Ibuprofen	9 a.m.	
Acetaminophen	12 p.m.	
Ibuprofen	3 p.m.	
Acetaminophen	6 p.m.	
Ibuprofen	9 p.m.	

MPEC rev :: MM/YY